
by boiling the state of the sta


Limonene is a liquid hydrocarbon found in orange peel. It can be extracted by boiling orange peel with water, using the apparatus shown below. The mixture of limonene all water distils at a temperature which is 1 °C below the boiling point of water.



(a) (i)	State the name of the piece of apparatus labelled A.	
	[1]	
(ii)	Suggest what the reading on the thermometer will be when the limonene-water mixture is being distilled.	
	°C [1]	
(iii)	Limonene is less dense than water. What information in the diagram shows this?	

A student used the pieces of equipment shown below to compare the concentration of alkali in two liquid oven cleaners.

Oven cleaners contain the alkali sodium hydroxide.



(a)	Name the pieces of equipment.
	c
	D[2
(b)	Outline how these pieces of equipment could be used in an investigation to compare the concentration of alkali in the two liquid oven cleaners.
	•


4 An aqueous solution **H** and a solid **J** were tested. The tests and some of the observations are in the following table. **H** was hydrochloric acid and **J** was a metal oxide.

Complete the observations in the table.

	tests	observations
(a)	A little of solution <b>H</b> was tested with Universal indicator paper.	colour
	The pH was recorded.	pH
(b)	To 1 cm <sup>3</sup> of solution <b>H</b> was added one spatula measure-of manganese(IV) oxide. The mixture was boiled. The gas was tested with damp blue litmus paper.	litmus paper turned pink, then white
(c)	To 1 cm <sup>3</sup> of solution <b>H</b> was added about 1 cm <sup>3</sup> of aqueous silver nitrate.	
(d)	To 1 cm <sup>3</sup> of solution <b>H</b> was added one spatula measure of sodium carbonate. The gas given off was tested with limewater.	
(e)	To 6 cm <sup>3</sup> of solution H was added solid J. The mixture was boiled gently. Two portions of the mixture were tested as follows.	
	(i) To 1 cm <sup>3</sup> of the mixture was added an excess of aqueous sodium hydroxide a little at a time.	white precipitate, dissolved in excess.
	(ii) To 1 cm <sup>3</sup> of the mixture was added an excess of aqueous ammonia a little at a time.	white precipitate, dissolved in excess.

	at a time.	white precipitate, dissolved in excess.
(f)	Name the gas given off in test (b).	
		[1]
(g)	Name the gas given off in test (d).	
		[1]
(h)	Identify the metal in oxide J.	
		[1]

The diagram shows apparatus used for strongly heating some magnesium.



(a) Name each piece of apparatus.

A	
В	
C	
D	[4

(b) Give two expected observations if the lid were removed during the heating.

1	
2	[2]

(d) Chromatography is used to identify simple carbohydrates, such as sugars, in plant material.




Fig. 5.2

A leaf is ground with 50% aqueous alcohol to give a colourless solution of the sugars. This solution is concentrated and a chromatogram is obtained. The paper is sprayed with resorcinol solution.

(i) A common use of ethanol is in alcoholic drinks. In this experiment it is used as a solvent. Give one other use.
[1]
(ii) Why is the datum line drawn in pencil?
[1]
iii) Suggest a reason why it is necessary to spray the chromatogram with resorcinol.
[2]
iv) Describe how chromatography could be used to show that the hydrolysis of starch produces only one sugar, glucose.
[2]

Chemistry is concerned with problem-solving and answering questions. For each of the following, describe any experiments that are needed and, when necessary, indicate how the results should be used to solve the problem.

(a)	as a	kel sulphate-7-water exists as bright green crystals and anhydrous nickel sulphate a yellow powder. How could you show that the action of heat on the hydrated salt is a ersible reaction?
	••••	
	••••	[3]
(b)	chlo	w could you prove that the percentage of water in an aqueous solution of sodium oride is 74%?
	••••	[4]
(c)	A s filte	mall piece of cement from a wall is dissolved in nitric acid. The resulting mixture is red to give a solution.
	(i)	What observation shows that the cement contains carbonate ions and what test is needed to confirm this?
		observation
		test
		result
	(ii)	How could you show that the solution contains calcium ions?
		reagent
		result
		reagent
		result
		[6]
(b)	is c	formula of an acid is either of the type HY or H <sub>2</sub> Y. How could you find out which one prrect by using aqueous solutions of the acid and of sodium hydroxide. Assume that a solutions have the same concentration, 0.1 mol/dm <sup>3</sup> .
	*****	
	••••	
		[1]

# Core 1

- a(i) (Liebig) condenser
- (ii) 99
- (iii) limonene floats on water / on top of the water

- a C = burette
  - D = pipette
- b five points from these

known volume of / pipette oven cleaner into flask / beaker named acid in burette use of indicator note volume of acid to neutralise oven cleaner repeat with other oven cleaner

- a colour red pH 1-3
- c white precipitate
- d fizz / effervescence / bubbles limewater gives a milky / white precipitate
- f chlorine
- g carbon dioxide
- h zinc

- A Bunsen burner а
  - B Tripod C Gauze D Crucible
- indication of burning, light / flame / glow white smoke / ash b

- i fuel or making esters or antiseptic or ethanoic acid or vinegar or thermometers
- ii does not dissolve or does not contain dyes
- iii two of these

to develop it or locating agent samples are colourless to make them visible

iv any two of these only one spot same position or Rf value compare with glucose

- heat green crystals
  add water to yellow powder
  goes green
- weigh solution
  evaporate to dryness
  weigh (solid)
  solution solid is water or mass of water / mass of solution \* 100
- c bubbles limewater goes cloudy

add sodium hydroxide white precipitate

add aqueous ammonia no precipitate or faint one

or flame test with red / orange / brick red flame

d any four of these indicator titration

find volume of sodium hydroxide to neutralise if volumes the same then HY

if twice volume of alkali then H<sub>2</sub>Y